

## Genotyping of *Toxoplasma gondii* Isolates with 15 Microsatellite Markers in a Single Multiplex PCR Assay<sup>∇</sup>

Daniel Ajzenberg,<sup>1,2\*</sup> Frédéric Collinet,<sup>2</sup> Aurélien Mercier,<sup>2</sup>  
Philippe Vignoles,<sup>2</sup> and Marie-Laure Dardé<sup>1,2</sup>

Centre National de Référence (CNR) *Toxoplasme*/Toxoplasma Biological Resource Center (BRC), Centre Hospitalier-Universitaire Dupuytren, Limoges, 87042, France,<sup>1</sup> and Laboratoire de Parasitologie-Mycologie, EA 3174-NETEC, Faculté de Médecine, Université de Limoges, Limoges, 87025, France<sup>2</sup>

Received 7 June 2010/Returned for modification 28 July 2010/Accepted 17 September 2010

**We developed an easy-to-use method for genotyping *Toxoplasma gondii* isolates in a single multiplex PCR assay with 15 microsatellite markers. This method was validated by testing 26 reference isolates that had been characterized with other sets of markers.**

The commonly used multilocus methods for genotyping *Toxoplasma gondii* strains employ PCR-restriction fragment length polymorphism (PCR-RFLP) analysis at 10 markers (16), sequencing at 8 introns of 5 loci (14), and length polymorphism analysis of 5 microsatellite (MS) markers (4). Advantages and disadvantages of these different markers have been reviewed elsewhere (16). Based on our experience within the *Toxoplasma* Biological Resource Center (BRC ToxoBS) and the National Reference Center for Toxoplasmosis, it appears that screening of a large number of clinical isolates for *T. gondii* strain typing should rely on an easy-to-use method with two different levels of discrimination. The first step of discrimination is the performed at the typing level: that is, employing the ability of markers to distinguish the major clonal lineages from atypical strains. In areas with a marked clonal population structure such as Europe, the genetic screening of clinical isolates should rapidly identify basic type II or III strains, which represent the majority of isolates, and atypical strains, which are the exception to the rule (3, 11). In this context, the identification of atypical strains is of clinical and epidemiological importance because atypical strains are usually associated with severe disease outcomes and contamination of individuals by non-European strains either during residence abroad or after consumption of imported meat (5, 9, 10). The second level of discrimination is the fingerprinting level, representing a high degree of discriminatory power for differentiating closely related strains belonging to the same lineage. This high-resolution analysis is required for identifying laboratory contaminations for diagnosis issues and for establishing a common source of infection among different infected individuals in an outbreak (7, 8).

Microsatellite (MS) sequences are tandem repeats of short (1 to 6 bp) DNA motifs that are ubiquitous in eukaryotic

genomes and undergo length changes due to insertion or deletion of one or multiple repeat units. The most commonly proposed mutation mechanism for MS sequences is strand slippage, occurring predominantly during replication (13). The numbers of repeat motifs differ in a population, thereby creating multiple alleles at an MS locus. MS loci are amplified by PCR using fluorescently labeled forward and unlabeled reverse primers. The dye-labeled products are separated by size using automated electrophoresis and identified by fluorescence detection. We previously designed a multiplex PCR assay with 5 MS markers that were able to reach the typing level but not the fingerprinting level of discrimination between strains (4). Here we developed an easy-to-use and rapid genotyping method which aimed to ensure both levels of genetic discrimination of *T. gondii* isolates in a single multiplex PCR assay using 15 MS markers located on 11 different chromosomes of *T. gondii*. In MS sequences with dinucleotide motifs, those with [TG/AC]<sub>n</sub> and [TC/AG]<sub>n</sub> repeats are known to exhibit lower mutability than those with [TA/AT]<sub>n</sub> repeats (13). Eight MS markers with a repeat motif consisting of [TG/AC]<sub>n</sub> or [TC/AG]<sub>n</sub> were used as typing markers: five markers (*TUB2*, *W35*, *TgM-A*, *B18*, and *B17*) were described in our previous report of a multiplex assay (4), *M33* was developed by others (6), and two markers (*IV.1* and *XI.1*) were designed for this study. Seven MS markers with a repeat motif consisting of [TA/AT]<sub>n</sub> were used as fingerprinting markers: five markers (*N60*, *N82*, *AA*, *N61*, and *N83*) were described by us in a previous study (2) whereas *M48* and *M102* were developed by others (6). For some markers, primers sequences were redesigned to allow PCR amplification and allele size discrimination in the multiplex assay. Chromosome locations and primer sequences for the 15 MS markers are listed in Table 1. Primers were synthesized by Applied Biosystems (Courtabœuf, France). PCR was carried out in a 25- $\mu$ l reaction mixture consisting of 12.5  $\mu$ l of 2 $\times$  Qiagen multiplex PCR Master Mix (Qiagen, France) and 5 pmol each primer. The volumes of DNA template were 1  $\mu$ l for DNA extracted from tachyzoites in cell culture or mouse ascitic fluid, 3  $\mu$ l for DNA extracted from bradyzoites in mouse brain, and 5  $\mu$ l for DNA extracted directly from clinical samples. Cycling conditions

\* Corresponding author. Mailing address: Centre National de Référence (CNR) *Toxoplasme*, Laboratoire de Parasitologie et de Mycologie, Centre Hospitalier-Universitaire Dupuytren, 2 avenue Martin Luther King, 87042 Limoges, France. Phone: 33555056160. Fax: 33555056177. E-mail: ajz@unilim.fr.

<sup>∇</sup> Published ahead of print on 29 September 2010.

TABLE 1. Microsatellite markers and PCR primers used for the multiplex PCR assay

Marker	Chromosome (position) <sup>a</sup>	Repeat motif(s)	Primer sequence <sup>b</sup>	Size range (bp)
<i>TUB2</i>	IX (974608 to 974896)	[TG/AC] <sub>n</sub>	(F) 5' 6-FAM-GTCCGGGTGTTTCCTACAAAA 3' (R) 5' TTGGCCAAAGACGAAGTTGT 3'	287–291
<i>W35</i>	II (633241 to 633482)	[TC/AG] <sub>n</sub> , [TG/AC] <sub>n</sub>	(F) 5' HEX-GGTTCACTGGATCTTCTCCAA 3' (R) 5' AATGAACGTCGCTTGTTC 3'	242–248
<i>TgM-A</i>	X (4824879 to 4825083)	[TG/AC] <sub>n</sub>	(F) 5' HEX-GGCGTCGACATGAGTTTCTC 3' (R) 5' TGGGCATGTAATGTAGAGATG 3'	203–211
<i>B18</i>	VIIa (2921536 to 2921693)	[TG/AC] <sub>n</sub>	(F) 5' 6-FAM-TGGTCTTACCCTTTCATCC 3' (R) 5' AGGGATAAGTTTCTTACAACGA 3'	156–170
<i>B17</i>	XII (6474746 to 6475079)	[TC/AG] <sub>n</sub>	(F) 5' HEX-AACAGACACCCGATGCCTAC 3' (R) 5' GGCAACAGGAGGTAGAGGAG 3'	334–366
<i>M33</i>	IV (672591 to 672760)	[TC/AG] <sub>n</sub>	(F) 5' 6-FAM-TACGCTTCGATTGTACCAG 3' (R) 5' TCTTTTCTCCCTTCGCTCT 3'	165–173
<i>IV.1</i>	IV (742419 to 742693)	[TG/AC] <sub>n</sub>	(F) 5' HEX-GAAGTTCGGCCTGTTCCCTC 3' (R) 5' TCTGCCTGAAAAGGAAAAGA 3'	272–282
<i>XI.1</i>	XI (189702 to 190058)	[TG/AC] <sub>n</sub>	(F) 5' 6-FAM-GCGTGTGACGAGTTCTGAAA 3' (R) 5' AAGTCCCCTGAAAAGCCAAT 3'	354–362
<i>M48</i>	Ia (332951 to 333166)	[TA/AT] <sub>n</sub>	(F) 5' 6-FAM-AACATGTGCGGTAAGATTTCG 3' (R) 5' CTCTTCACTGAGCGCCTTTC 3'	209–243
<i>M102</i>	VIIa (3093491 to 3093664)	[TA/AT] <sub>n</sub>	(F) 5' NED-CAGTCCAGGCATACCTCACC 3' (R) 5' CAATCCCAAAATCCCAAAACC 3'	164–196
<i>N60</i>	Ib (1766079 to 1766221)	[TA/AT] <sub>n</sub>	(F) 5' NED-GAATCGTCGAGGTGCTATCC 3' (R) 5' AACGGTTGACCTGTGGCGAGT 3'	132–157
<i>N82</i>	XII (1621472 to 1621585)	[TA/AT] <sub>n</sub>	(F) 5' HEX-TGCGTGCTTGTGAGAGTTC 3' (R) 5' GCGTCCTTGACATGCACAT 3'	105–145
<i>AA</i>	VIII (5836880 to 5837144)	[TA/AT] <sub>n</sub>	(F) 5' NED-GATGTCCGGTCAATTTTGTCT 3' (R) 5' GACGGGAAGGACAGAAACAC 3'	251–332
<i>N61</i>	VIIb (4217145 to 4217238)	[TA/AT] <sub>n</sub>	(F) 5' 6-FAM-ATCGGCGGTGGTTGTAGAT 3' (R) 5' CCTGATGTTGATGTAAGGATGC 3'	79–123
<i>N83</i>	X (1772898 to 1773209)	[TA/AT] <sub>n</sub>	(F) 5' 6-FAM-ATGGGTGAACAGCGTAGACA 3' (R) 5' GCAGGACGAAGAGGATGAGA 3'	306–338

<sup>a</sup> Based on position within each chromosome of strain ME49 as given in the Toxo DB website (<http://toxodb.org>).

<sup>b</sup> (F), forward primer; (R), reverse primer. In each pair of primers, the forward primer was labeled at the 5' end with fluorescein: 6-carboxyfluorescein (6-FAM) for MS *TUB2*, *XI.1*, *B18*, *N83*, *N61*, *M33*, and *M48*, hexachlorofluorescein (HEX) for MS *TgM-A*, *B17*, *N82*, *W35*, and *IV.1*, and 2',7',8'-benzo-5'-fluoro-2',4,7-trichloro-5-carboxyfluorescein (NED) for MS *AA*, *N60*, and *M102*.

were 15 min at 95°C; 30 s at 94°C, 3 min at 61°C, and 30 s at 72°C (35 cycles); and 30 min at 60°C. PCR products were diluted in deionized formamide as follows: 1/20 for DNA extracted from tachyzoites in cell culture or mouse ascitic fluid, 1/10 for DNA extracted from bradyzoites in mouse brain, and 1/2 for DNA extracted directly from clinical samples. One microliter of each diluted PCR product was mixed with 0.5 µl of a dye-labeled size standard (ROX 500; Applied Biosystems) and 23.5 µl of deionized formamide. This mixture was denatured at 95°C for 5 min and then electrophoresed using an automatic sequencer (ABI PRISM 3130xl; Applied Biosystems). The sizes of the alleles in base pairs were esti-

mated using GeneMapper analysis software (version 4.0; Applied Biosystems) (Fig. 1).

We used the 15-MS multiplex assay to genotype 26 reference isolates that had been extensively characterized with other sets of markers. These strains have been classified either into the three main clonal lineages (types I, II, and III as classified by PCR-RFLP markers, corresponding to haplogroups 1, 2, and 3 as classified by intron sequencing, respectively) or into atypical genotypes by PCR-RFLP markers corresponding to haplogroups 4 to 11 by intron sequencing (Table 2). The genotyping results of the 15-MS multiplex assay showed that the 8 typing markers (*TUB2*, *W35*, *TgM-A*, *B18*, *B17*, *M33*, *IV.1*, and *XI.1*)




FIG. 1. Electrophoresis of the 15-microsatellite multiplex PCR amplification products from strain P89 by the use of an automatic sequencer and GeneMapper software. The x axis indicates a size fragment scale (in base pairs). The y axis indicates a peak height scale corresponding to the fluorescence intensities of detected peaks.

could readily distinguish clonal types I, II, and III from all the atypical genotypes (Table 2). In each of clonal lineages I to III or haplogroups 1 to 3, representative strains had uniform typing patterns, except for one for lineage II, where allele 244 at marker *W35* was identified for isolate BOU. Such variants within clonal lineage II have also been previously determined at the *Apico* locus with PCR-RFLP markers (17). Atypical isolates were characterized by a different mix of alleles of haplogroups 1 to 3 and by additional atypical alleles at some loci. The detection of atypical alleles by the MS markers underscores the ability of these markers to capture the genetic diversity of highly divergent strains such as South American strains (1, 14, 15). The 7 fingerprinting markers (*M48*, *M102*, *N60*, *N82*, *AA*, *N61*, and *N83*) provided enhanced genetic resolution, beyond the typing level, in distinguishing closely related isolates within one clonal lineage or haplogroup (Table 2). The Simpson's index of diversity ( $D$ ) (2) was maximal ( $D = 1$ ), since all reference isolates had a unique genotype as classified with the 15 MS markers.

The association of the 15 MS markers with different mutability patterns enabled two different levels of genetic resolution for differentiating *T. gondii* strains at the typing level (types I, II, and III versus atypical strains) and at the fingerprinting level (closely related isolates within a clonal lineage). Multilocus sequence-based typing is the best tool for estimating the true rate of DNA polymorphism in phylogenetic studies, but the

increased cost and investment of time makes that method unsuitable for genetic screening of a large number of isolates. PCR-RFLP markers are preferred tools for the latter purpose, but their low resolution level makes them inadequate for distinguishing genetically closely related strains. The 15-MS multiplex assay is therefore the best available tool to reveal whether two isolates are genetically identical or different, i.e., to identify a common source of infection in an outbreak, a laboratory contamination, and mixed infections. The limitations of this assay are the requirement for the availability of an automated sequencer and the sensitivity of the method, which has been estimated by testing serial dilutions of RH strain tachyzoites to be between 50 to 100 *T. gondii* genome equivalents per 5  $\mu$ l of DNA sample. Testing this method on samples with lower DNA concentrations is not recommended because of incomplete amplification of markers or difficulties in interpreting the results due to overlaps of microsatellite peaks with nonspecific PCR products. Compared with other genotyping methods, the sensitivity appears similar to that reported with intron sequencing. PCR-RFLP analysis can detect as few as 10 *T. gondii* genome equivalents per PCR, but such a threshold of detection is reached only by nested PCR assays (16). We did not develop a nested PCR assay for improving the sensitivity of our 15-MS multiplex assay since, in our experience, double-round PCR amplification using this method represents too

TABLE 2. Genotyping results of 26 reference strains with 15 MS markers in a single multiplex PCR assay

Type (haplogroup) <sup>a</sup>	Isolate <sup>b</sup>	Origin (host) <sup>c</sup>	Microsatellite marker														
			TUB2	W35	TgM-A	B18	B17	M33	IV.1	XI.1	M48	M102	N60	N82	AA	N61	N83
I (1)	P	France (H)	291	248	209	160	342	169	274	358	209	166	145	121	267	87	308
I (1)	CT1	USA (cow)	291	248	209	160	342	169	274	358	209	168	145	119	265	87	306
I (1)	GIL	France (H)	291	248	209	160	342	169	274	358	209	166	147	119	265	87	306
I (1)	BK	Netherlands (H)	291	248	209	160	342	169	274	358	209	166	145	119	265	87	316
II (2)	NTE	Germany (H)	289	242	207	158	336	169	274	356	221	176	140	113	269	103	310
II (2)	PTG	USA (sheep)	289	242	207	158	336	169	274	356	215	174	142	111	265	91	310
II (2)	BOU	France (H)	289	244	207	158	336	169	274	356	213	178	140	113	259	99	310
III (3)	CTG	USA (cat)	289	242	205	160	336	165	278	356	215	190	147	111	269	89	312
III (3)	NED	France (H)	289	242	205	160	336	165	278	356	209	190	147	111	267	91	312
III (3)	M7741	USA (sheep)	289	242	205	160	336	165	278	356	215	190	147	111	267	91	312
III (3)	VEG	USA (H)	289	242	205	160	336	165	278	356	213	188	153	111	267	89	312
Atypical (4)	MAS	France (H)	291	242	205	162	362	169	272	358	221	166	142	111	332	95	338
Atypical (4)	CASTELLS	Uruguay (sheep)	287	242	207	158	358	169	274	356	239	164	138	109	283	87	324
Atypical (4)	TgCatBr1	Brazil (cat)	289	242	205	160	342	165	278	358	233	164	147	111	316	89	308
Atypical (5)	GUY-KOE	Fr. Guiana (H)	289	246	203	160	337	165	274	356	209	172	136	111	251	109	310
Atypical (5)	GUY-MAT	Fr. Guiana (H)	291	242	203	160	339	165	272	358	221	174	138	107	277	95	312
Atypical (5)	RUB	Fr. Guiana (H)	289	242	205	170	360	167	274	356	223	190	142	109	259	85	312
Atypical (6)	GPHT	France (H)	291	248	205	160	342	165	274	354	229	166	147	111	283	91	306
Atypical (6)	BOF	Belgium (H)	291	248	205	160	342	165	274	354	227	166	147	111	273	89	306
Atypical (7)	CAST	USA (H)	291	242	205	158	342	167	276	356	211	168	147	119	279	87	306
Atypical (8)	TgCatBr5	Brazil (cat)	291	242	205	160	362	165	278	356	237	174	140	111	265	89	314
Atypical (9)	P89	USA (pig)	291	242	205	160	348	165	278	356	213	190	142	111	261	87	314
Atypical (9)	TgCatBr3	Brazil (cat)	289	242	205	160	348	165	278	356	213	190	142	111	263	113	312
Atypical (10)	VAND	Fr. Guiana (H)	291	242	203	162	344	167	276	356	217	170	142	113	277	91	308
Atypical (10)	GUY-DOS	Fr. Guiana (H)	289	246	203	160	344	167	272	356	229	176	142	113	263	85	312
Atypical (11)	COUGAR	Canada (cougar)	289	242	205	158	336	169	274	354	219	174	151	119	259	79	332

<sup>a</sup> The types were defined by analysis of single nucleotide polymorphisms at 5 or 10 PCR-RFLP markers (12, 17); the haplogroups were defined by analysis of sequence polymorphisms within 8 introns of five unlinked loci (14).

<sup>b</sup> Strain P is also known as strain PT; strain PTG is a clone of the ME49 strain; strain CTG is also known as strain CEP or C; strains GUY-KOE, GUY-MAT, and GUY-DOS are also known as strains GUY-2002-KOE, GUY-2002-MAT, and GUY-2001-DOS, respectively; strain P89 is also known as strain TgPgUs15; strain COUGAR is also known as strain TgCgCa1 or COUG.

<sup>c</sup> Fr. Guiana, French Guiana; H, human; UK, United Kingdom; USA, United States.

high a risk of PCR contamination and often leads to misleading results.

We thank Jitender Dubey, David Sibley, and Chunlei Su for kindly providing DNA samples from reference strains. We are indebted to the following members of the network of the *Toxoplasma* Biological Resource Center for providing *T. gondii* isolates from human cases of toxoplasmosis: Sahar Albaba, Dominique Aubert, Jacques Berthonneau, Marie-Hélène Bessières, Rachida Boukhari, Marie-Pierre Brenier-Pinchart, Bernard Carme, Sophie Cassaing, Bernard Cimon, Bernard Couprie, Bernadette Cuisenier, Laurence Delhaes, Magalie Demar, Nicole Desbois, Chantal Duhamel, Than-Hai Duong, Loïc Favennec, Denis Filisetti, Pierre Flori, Jacqueline Franck, Héléne Fricker-Hidalgo, Françoise Gay-Andrieu, Catherine Kauffmann-Lacroix, Marie Machouart, Pierre Marty, Jean Menotti, Gilles Nevez, Luc Paris, Hervé Pelloux, Christelle Pomares, Francine Pratlong, Dorothée Quinio, Florence Robert-Gangneux, Philippe Thulliez, Anne Totet, Nathalie Van Langendonck, Isabelle Villena, and Héléne Yera.

#### REFERENCES

- Ajzenberg, D., A. L. Banuls, C. Su, A. Dumètre, M. Demar, B. Carme, and M. L. Dardé. 2004. Genetic diversity, clonality and sexuality in *Toxoplasma gondii*. *Int. J. Parasitol.* **34**:1185–1196.
- Ajzenberg, D., A. L. Banuls, M. Tibayrenc, and M. L. Dardé. 2002. Microsatellite analysis of *Toxoplasma gondii* shows considerable polymorphism structured into two main clonal groups. *Int. J. Parasitol.* **32**:27–38.
- Ajzenberg, D., N. Cogné, L. Paris, M. H. Bessières, P. Thulliez, D. Filisetti, H. Pelloux, P. Marty, and M. L. Dardé. 2002. Genotype of 86 *Toxoplasma gondii* isolates associated with human congenital toxoplasmosis, and correlation with clinical findings. *J. Infect. Dis.* **186**:684–689.
- Ajzenberg, D., A. Dumètre, and M. L. Dardé. 2005. Multiplex PCR for typing strains of *Toxoplasma gondii*. *J. Clin. Microbiol.* **43**:1940–1943.
- Ajzenberg, D., H. Yera, P. Marty, L. Paris, F. Dalle, J. Menotti, D. Aubert, J. Franck, M. H. Bessières, D. Quinio, H. Pelloux, L. Delhaes, N. Desbois, P. Thulliez, F. Robert-Gangneux, C. Kauffmann-Lacroix, S. Pujol, M. Rabodonirina, M. E. Bougnoux, B. Cuisenier, C. Duhamel, T. H. Duong, D. Filisetti, P. Flori, F. Gay-Andrieu, F. Pratlong, G. Nevez, A. Totet, B. Carme, H. Bonnahau, M. L. Dardé, and I. Villena. 2009. Genotype of 88 *Toxoplasma gondii* isolates associated with toxoplasmosis in immunocompromised patients and correlation with clinical findings. *J. Infect. Dis.* **199**:1155–1167.
- Blackston, C. R., J. P. Dubey, E. Dotson, C. Su, P. Thulliez, D. Sibley, and T. Lehmann. 2001. High-resolution typing of *Toxoplasma gondii* using microsatellite loci. *J. Parasitol.* **87**:1472–1475.
- Carme, B., D. Ajzenberg, M. Demar, S. Simon, M. L. Dardé, B. Maubert, and B. de Thoisy. 2009. Outbreaks of toxoplasmosis in a captive breeding colony of squirrel monkeys. *Vet. Parasitol.* **163**:132–135.
- Demar, M., D. Ajzenberg, D. Maubon, F. Djossou, D. Panchoe, W. Punwasi, N. Valery, C. Peneau, J. L. Daigre, C. Aznar, B. Cottrelle, L. Terzan, M. L. Dardé, and B. Carme. 2007. Fatal outbreak of human toxoplasmosis along the Maroni River: epidemiological, clinical, and parasitological aspects. *Clin. Infect. Dis.* **45**:e88–e95.
- De Salvador-Guillouët, F., D. Ajzenberg, S. Chaillou-Opitz, M. C. Saint-Paul, B. Dunais, P. Dellamonica, and P. Marty. 2006. Severe pneumonia during primary infection with an atypical strain of *Toxoplasma gondii* in an immunocompetent young man. *J. Infect.* **53**:e47–e50.
- Elbez-Rubinstein, A., D. Ajzenberg, M. L. Dardé, R. Cohen, A. Dumètre, H. Yera, E. Gondon, J. C. Janaud, and P. Thulliez. 2009. Congenital toxoplasmosis and reinfection during pregnancy: case report, strain characterization, experimental model of reinfection, and review. *J. Infect. Dis.* **199**:280–285.
- Halos, L., A. Thebault, D. Aubert, M. Thomas, C. Perret, R. Geers, A. Alliot, S. Escotte-Binet, D. Ajzenberg, M. L. Dardé, B. Durand, P. Boireau, and I. Villena. 2010. An innovative survey underlining the significant level of contamination by *Toxoplasma gondii* of ovine meat consumed in France. *Int. J. Parasitol.* **40**:193–200.
- Howe, D. K., and L. D. Sibley. 1995. *Toxoplasma gondii* comprises three clonal lineages: correlation of parasite genotype with human disease. *J. Infect. Dis.* **172**:1561–1566.
- Kelkar, Y. D., S. Tyekucheva, F. Chiaromonte, and K. D. Makova. 2008. The genome-wide determinants of human and chimpanzee microsatellite evolution. *Genome Res.* **18**:30–38.
- Khan, A., B. Fux, C. Su, J. P. Dubey, M. L. Dardé, J. W. Ajioka, B. M.

- Rosenthal, and L. D. Sibley.** 2007. Recent transcontinental sweep of *Toxoplasma gondii* driven by a single monomorphic chromosome. Proc. Natl. Acad. Sci. U. S. A. **104**:14872–14877.
15. **Pena, H. F., S. M. Gennari, J. P. Dubey, and C. Su.** 2008. Population structure and mouse-virulence of *Toxoplasma gondii* in Brazil. Int. J. Parasitol. **38**:561–569.
16. **Su, C., E. K. Shwab, P. Zhou, X. Q. Zhu, and J. P. Dubey.** 2010. Moving towards an integrated approach to molecular detection and identification of *Toxoplasma gondii*. Parasitology **137**:1–11.
17. **Su, C., X. Zhang, and J. P. Dubey.** 2006. Genotyping of *Toxoplasma gondii* by multilocus PCR-RFLP markers: a high resolution and simple method for identification of parasites. Int. J. Parasitol. **36**:841–848.